



EDITORIAL

Reducing the global cancer burden with gastrointestinal screening: China's 30 years practice

Lei Yang^{1,2*}, Li Feng^{1*}, Yong Zhu^{1*}, Ning Wang², Xinpu Lu³, Fanghui Gu³, Xiaotian Zhang^{1,4,5}, Jiafu Ji^{4,5}

¹Peking University Cancer Hospital (Inner Mongolia Campus)/Affiliated Cancer Hospital of Inner Mongolia Medical University, Inner Mongolia Cancer Center, Hohhot 010020, China; ²Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education/Beijing), Beijing Office for Cancer Prevention and Control, Peking University Cancer Hospital & Institute, Beijing 100142, China; ³Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education), Peking University Cancer Hospital & Institute, Beijing 100142, China; ⁴Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education/Beijing), Department of Gastrointestinal Oncology, Peking University Cancer Hospital & Institute, Beijing 100142, China; ⁵State Key Laboratory of Holistic Integrative Management of Gastrointestinal Cancers, Beijing Key Laboratory of Carcinogenesis and Translational Research, Gastrointestinal Cancer Center, Peking University Cancer Hospital & Institute, Beijing 100142, China

Gastrointestinal (GI) cancers include esophageal [EC (International Classification of Disease 10th revision: C15)], gastric [GC (C16)], and colorectal [CRC (C18-C21)] cancers. China is a high-incidence country of GI cancers. According to GLOBOCAN2020, the number of new cases and deaths from GI cancers in China was 1,368,407 and 961,086, respectively, accounting for approximately 38% and 43% of the global new cases and deaths, respectively, in 2020¹. Upper GI cancers, including EC and GC, dominate in developing countries, while CRC is prevalent in developed countries².

GI cancers derived from distinct but associated origins have diverse clinical features yet share some similar characteristics. Esophageal squamous cell carcinoma (ESCC) is the most common pathologic EC subtype in China. The main risk factors for ESCC include a preference for eating hot foods, cigarette smoking, alcohol consumption, a high-salt diet, and bad oral hygiene³. For GC, especially non-cardia GC, the main

risk factors are *Helicobacter pylori* infection, a high-salt diet, insufficient fruit and vegetable intake, cigarette smoking, alcohol consumption⁴, and a family history of GC⁵. Consequently, upper GI cancers, particularly ESCC and non-cardia GC, are prevalent in countries with a low human development index (HDI), especially China. At the same time, CRC is more common in regions with a high HDI, with the main risk factors including the consumption of processed meats, overweight and obesity, lack of physical activity, and a family history of CRC⁶.

As global urbanization accelerates and economies develop rapidly, countries undergoing economic transitions have experienced notable shifts in digestive tract cancer patterns². In China, this shift has manifested as a declining incidence of upper GI cancers, which is mainly due to the decreasing proportion of infectious factors. Conversely, CRC, which is primarily linked to a westernized lifestyle⁷, has been reported to have an upward trend. Recent data on cancer mortality in China revealed significant decreases in mortality and years of life lost (YLL) rates for all GI cancers in urban areas after age-standardization, except the CRC mortality in males⁸.

Owing to the large and aging population in China, upper GI cancers, despite a decreasing mortality rate, claimed 464,490 lives in 2020, resulting in 9,916,570 YLLs. CRC, with increasing annual incidence and mortality rates, causes 182,430 deaths annually, resulting in 3,884,250 YLLs⁸. The prevention and control of GI cancers are exceptionally serious matters in China.

*These authors contributed equally to this work.

Correspondence to: Xiaotian Zhang and Jiafu Ji

E-mail: zhangxiaotian@bjmu.edu.cn and jijiafu@hsc.pku.edu.cn

ORCID ID: <https://orcid.org/0000-0001-6936-2622>

and <https://orcid.org/0000-0001-6878-5543>

Received January 25, 2024; accepted February 22, 2024.

Available at www.cancerbiomed.org

©2024 Cancer Biology & Medicine. Creative Commons

Attribution-NonCommercial 4.0 International License




Figure 1 Key milestones for gastrointestinal cancer screening in China.

Research in regions with a high global incidence of upper GI cancers indicated that early screening can reduce the incidence and mortality. Moreover, evidence from CRC screening in the United States suggests that 50% of the decline in the mortality rate of CRC is due to early diagnosis and treatment⁹.

Over the past few decades, China has implemented a series of cancer prevention and control measures that have somewhat reduced the global burden of GI cancers. The Cancer Prevention and Control Outline in China (2004–2010), which was released in 2003 by the Chinese government¹⁰, initiated the following three large-scale national population-based cancer screening programs: the Cancer Screening Program in Rural Areas, which was initiated in 2005; the Cancer Screening Program in Huai River Areas, which was initiated in 2007; and the Cervical Cancer and Breast Cancer Screening Program for Women in Rural Areas, which was initiated in 2009¹¹. With the subsequent release of the Plan for Prevention and Control of Chronic Disease in China (2012–2015) in 2012¹², the Cancer Screening Program in Urban China (CanSPUC) was launched, which focused on non-high-incidence areas for cancer screening. These four national early cancer diagnosis and treatment programs laid the foundation for large-scale population-based cancer screening projects in China and are currently ongoing. Guided by the Healthy China 2030¹³, the Medium-to-Long-Term Plan for Prevention and Control of Chronic Diseases (2017–2025), and the Plan for Cancer Prevention and Control (2019–2022)¹¹ were released. In the most recently released Action Plan for Cancer Prevention and Control

(2023–2030)¹⁴, China announced strengthening the capacity of cancer screening and early diagnosis and set a key target to improve the overall 5-year cancer survival to 46.6% by 2030 (Figure 1). Indeed, China has entered a new era in cancer prevention and control. This article summarizes 30 years of GI cancer screening and early detection practices in China, discussing the disease burden trends, to guide the formulation of cancer screening strategies in China and other developing countries.

EC

Epidemiology and disease burden

China, a high-risk country for EC, accounted for approximately 55% of global EC cases in 2020. EC is the 5th most common cause of cancer-related deaths and the 6th most prevalent cancer type in China. The incidence of EC in males is 1.6-fold higher than females. From 2000–2016, the annual percentage change (APC) in the EC incidence decreased by 3.9% in males and 6.4% in females, while the mortality rate decreased by 4.1% in males and 6.3% in females¹⁵. This downward trend was more pronounced among females. EC is comprised of two main pathologic types [ESCC and esophageal adenocarcinoma (EAC)]. EAC is common in western countries, whereas ESCC accounts for > 90% of EC cases and is more prevalent in east asian countries, particularly in China¹⁶. The Henan, Hebei, and Shanxi provinces around the Taihang Mountains are the high-risk regions for EC in China¹⁷.

In economically developed areas like Beijing, EC ranks ninth among male cancer cases but does not enter the top ten for females. The age-standardized incidence rate of EC in Beijing is 3.59 per 100,000, lower than the national average of 11.13 per 100,000¹⁸. Currently, the 5-year relative survival of EC in China is 30.3%¹⁹, but early-detected EC can exceed 90%²⁰. Detecting EC at an earlier and potentially curable stage is crucial for improving patient survival.

EC screening in China

In Lin County, Henan Province, China has developed early detection methods for EC (esophageal balloon and sponge cytology) since the late 1960s¹¹. Regional EC screening programs that involve performing endoscopic screening for residents 40–69 years of age in rural high-risk areas were initiated in 2005. EC screening is still included in the subsequent Huai River program and CanSPUC¹¹. Over the past 30 years, China has established a screening modality for EC involving endoscopy with iodine staining and indicative biopsy for individuals with high risk and those with precursor lesions²¹. Research in high-risk rural areas has shown that endoscopic screening for precursor lesions, along with endoscopic resection or ablation of dysplastic lesions, reduces ESCC development and mortality²². In a 10-year cohort in Cixian County, Hebei Province, screening reduced EC cumulative incidence and mortality rates by 29.6% and 33.7%, respectively, compared to the control group²³. In Hua County, Henan Province, the screening group displayed a 47% and 66% reduction in cumulative EC incidence and mortality, respectively²⁴. However, the abovementioned research was based on cohort studies in areas with very a high incidence of EC. Evidence on the benefits of EC screening from randomized control trials are still lacking in China. Based on evidence from high-risk field studies spanning three decades, the current screening strategy in China prioritizes high-incidence areas or high-risk populations²⁵. In regions with limited endoscopic resources, accurate identification of high-risk individuals is crucial for EC screening. Unlike GC, which is an infection-associated malignancy, there are currently no definite high-risk factors for EC³. To enhance the benefits from ESCC screening, Li et al.²⁶ proposed a “two-step” precision screening modality in China. First, before baseline screening, a model evaluating individual risk for “currently carrying malignant lesions in the esophagus” was adopted

to identify high-risk subgroups. After the initial screening, a risk prediction model based on multiple factors, including the pathologic diagnosis and iodine staining status at baseline screening, was applied to evaluate the risk of progression of precursor lesions to malignant lesions. This protocol notably increased the protection rate of surveillance for severe dysplasia cases that had been missed with traditional pathology-based surveillance. One limitation of this study was that extrapolation of the results was restricted.

Nevertheless, the two-stage ESCC screening strategy effectively identified more EC patients while saving resources. As the number of individuals undergoing opportunistic screening increases²⁷, this strategy can also be applied to follow-up monitoring and management of those who undergo baseline EC screening. Despite the rapid development of liquid biopsy and multi-omics technologies, there are currently no recommended biomarkers for EC screening²⁸.

GC

Epidemiology and disease burden

The incidence and mortality rates of GC rank 3rd among all cancer types in China, with an estimated male-to-female incidence ratio of approximately 2:1. The annual percentage change for the incidence of GC in China decreased by 3.0% in males and 2.9% in females, whereas the mortality rate decreased by 3.4% in males and 4.2% in females from 2000–2016¹⁵. GC has been divided anatomically into two main subsites: cardia GC, arising from the gastroesophageal junction; and non-cardia GC, arising from more distal regions of the stomach (also referred to as distal GC)²⁹. Distal GC is more common in the Chinese population, for which *H. pylori* infection is a main risk factor. In recent decades, the incidence of distal GC has decreased, likely due to improved hygiene standards, increased intake of fresh fruits and vegetables, reduced consumption of salt-preserved foods, and *H. pylori* eradication³⁰. The age-standardized rate (ASR) for GC mortality was 40.8 per 100,000 in males and 18.6 per 100,000 in females between 1990 and 1992³¹. The ASR for GC mortality decreased to 17.77 per 100,000 in males and 7.13 per 100,000 in females in 2016¹⁵ and is expected to continue to decline in the future. Currently, the 5-year relative survival for GC in China is 35.1%¹⁹. Japan and South Korea are also high-incidence countries for GC, but report a much higher

5-year survival due to a higher proportion of early-stage diagnosis³².

GC screening in China

In China, scientific pilot study related to GC screening lags behind that of EC screening. The Peking University Cancer Hospital initiated a pilot population-based program for the early diagnosis and treatment of GC in Linqu County, Shandong Province, a high-incidence area in China, in 1983³³. The regional GC screening program began in 2005 in rural areas. Based on this program, three high-incidence areas for GC (Linqu County in Shandong Province, Zhuanghe County in Liaoning Province, and Wuwei County in Gansu Province) were selected as high-risk sites³⁴. The cancer screening program for GC in the Huai River was launched in 2007. In the subsequent CanSPUC, which was launched in 2012, GC screening were also included in the screening programs. The target population expanded from rural high-incidence areas to high-risk populations in urban areas where the risk was low-to-moderate. In the aforementioned National Polit GC Screening Programs in China, the questionnaire-based risk assessment tool was used as the first step to select high-risk individuals. The second step involved an endoscopic examination in a designated hospital. The baseline examination findings directed follow-up recommendations for each individual²¹. In a multicenter population-based cohort study in high-risk areas for upper GI cancer in China, one-time endoscopic screening reduced the mortality of non-cardia GC by 61% in males and 64% in females³⁵. Another population-based study in Linqu, China showed that receiving endoscopic screening significantly decreased the risk of invasive GC [age- and gender-adjusted relative risk (RR): 0.69, 95% confidence interval (CI): 0.52–0.92] and GC-specific mortality (RR: 0.33, 95% CI: 0.20–0.56), particularly for non-cardia GC³⁶. Unlike the risk factors for EC, *H. pylori*-induced gastritis is likely to be a major step in the development of GC, and therefore, serum markers associated with chronic atrophic gastritis and intestinal metaplasia are used as potential indicators for identifying high-risk individuals.

Consequently, population-based *H. pylori* “test and treat” strategies³⁷ were used in the high-risk field with a focus on serologic determination of five biomarkers (PGI, PGII, PGI:PGII ratio, anti-*H. pylori* antibodies, and gastrin-17)^{34,38}. Patients with different risks underwent various endoscopy

screening modalities based on risk stratification strategies²⁹. A meta-analysis of seven randomized controlled trials (RCTs) involving *H. pylori* eradication in healthy individuals demonstrated a 46% decrease in the incidence of GC. Among the four studies that assessed subsequent mortality, there was a significant decrease in cancer-specific mortality among treated individuals³⁹.

CRC

Epidemiology and disease burden

CRC is the 2nd most commonly diagnosed cancer and 4th leading cause of death in China¹⁵. A westernized lifestyle, which is characterized by excessive consumption of processed meats, overweight/obesity, and prolonged sedentary behavior, is a risk factor for the development of CRC². The annual percentage change in the incidence of CRC in China increased by 2.4% in males and 1.2% in females from 2000–2016, with a 1.3% increase in the mortality rate in males and no significant increase in the mortality rate in females¹⁵. The ranking of CRC deaths is positively correlated with the level of economic development. According to cancer mortality burden data from the Chinese Center for Disease Control and Prevention in 2020, CRC ranked as the 2nd leading cause of cancer-related deaths in males in Beijing and Shanghai, 3rd in Jiangsu and Guangzhou, and lower in other provinces. Among females, the ranking of deaths also exhibits similar characteristics to that of males across different economic development regions⁸.

Of adults aged 50–75 years, 58% underwent colonoscopy screening in the US in 2013, prompting a national campaign in 2014 that aimed to increase the CRC screening prevalence to 80% in the eligible population by 2018⁴⁰. Reaching this goal is estimated to avert 280,000 new cases and 200,000 deaths of CRC by 2030⁴¹. The latest data show that since the nationwide CRC screening recommendation in the United States, the mortality rate for CRC decreased by 55% among males and 61% among females⁴². The CRC screening experience in the United States warrants adoption in China.

CRC screening in China

Significant benefits can be obtained from CRC screening because grossly visible lesions typically take 10–15 years to

progress to CRC, leaving a wide time window for early diagnosis⁴³. Screening can remove precancerous lesions *via* colonoscopy or detect cancer at an earlier stage, thereby reducing the incidence and mortality, and alleviating the CRC burden⁴⁴. Adenomas and serrated polyps, the two primary subtypes of lesions, are precursors to most sporadic CRCs, 85%–90% of which evolve from adenomas². A sequential screening program combining high-risk factors and fecal occult blood testing (FOBT) has been implemented in the general population of Jiashan County, Zhejiang Province, China since 1989⁴⁵. The individual with a positive result will then undergo further colonoscopy screening. Fecal immunochemical testing (FIT) was not initially used as the primary screening method for high-risk individuals in the CanSPUC. Instead, those individuals identified as high risk based on questionnaire were referred to designated hospitals for colonoscopy screening⁴⁶. However, recognizing that the FIT can improve compliance with colonoscopy screening in high-risk populations⁴⁷, the FIT and questionnaires were included as the primary screening methods to identify high-risk individuals for CRC in CanSPUC beginning in 2016⁴⁸. Although FIT has been recommended as a first-step population screening approach in China, the accuracy of FIT is low. FIT-DNA is more sensitive for detecting advanced precancerous lesions and CRC than FIT, but has more false-positive results and is expensive⁴⁹. Consequently, FIT-DNA is not recommended for use in China's National Cancer Screening Programs. Although comprehensive CRC screening strategies have been implemented in European countries and in the United States for decades, China lacks a nationwide CRC screening plan. Furthermore, there is still no evidence from large-scale population screening in China demonstrating that colonoscopic screening for CRC can reduce mortality rates.

The best evidence to support the use of colonoscopy in China Guidelines for the Screening, Early Detection, and Early Treatment of Colorectal Cancer (2020, Beijing) was limited to meta-analysis results from 5 cohort studies, which estimated that colorectal screening was associated with a 56% decrease in the incidence of CRC and a 57% decrease in the risk of death⁵⁰. Most evidence released based on the National Screening Program in China always focus on the uptake and early detection rates of precancerous lesions and cancer⁴⁷. The National Cancer Center of China initiated a randomized controlled trial for CRC screening in 2019 but the results have not been published⁵¹.

Challenges for GI cancer screening

China has a vast territory and large population with significant variations in GI cancer disease burden and healthcare resources. National pilot cancer screening programs and population-based cohort studies launched over the past three decades have played an essential role in exploring suitable screening modalities in China¹¹. However, many barriers and challenges have not been addressed.

Identification of high-risk individuals

There are no nationwide screening recommendations for GI cancers in the general population in China⁵⁰. A two-step screening strategy (i.e., a non-invasive or minimally invasive approach to select high-risk individuals and those who should undergo endoscopy) for GI cancer screening in high-incidence areas is cost-effective. The accurate identification of high-risk individuals who may benefit the most from screening is crucial in low- and moderate-risk areas⁴⁷. Currently, high-risk population assessment methods used for specific age groups in national GI cancer screening programs, including questionnaires^{11,46}, risk prediction models³³, and blood-based biomarkers⁵⁰, all have limited sensitivity and positive predictive values. Moreover, the cost-effectiveness of these assessment methods in population settings remains unclear. New emerging technologies, such as multiomics combined with artificial intelligence for cancer screening, are still in the research stage. Some marketed products are expensive and have not reached a level for widespread use in population screening⁵².

Another critical issue is the low uptake rate in GI cancer screening due to the invasive nature of the screening procedures. Studies from one of the CanSPUC screening sites (Beijing) indicated that the uptake rates of GC and CRC screening was 15.9%⁵³ and 25.2%⁵⁴, respectively. The uptake rate of CRC screening in Beijing rural areas was higher than urban areas (58.9% *vs.* 25.2%)⁵⁵, possibly because rural areas have adopted a combined strategy of FIT and questionnaires for high-risk population assessment⁵⁶. However, the overall uptake rate of national digestive system cancer screening is low, and patient compliance directly affects screening effectiveness⁵⁷. To address the low uptake rates, it is essential to strengthen scientific health promotion campaigns to improve

awareness of the cancer screening benefits of individuals and seek alternative methods that can replace the existing invasive screening methods, making screening simple, non-invasive, and easily acceptable to examinees.

Endoscopy examination

The performance quality of GI cancer screening by endoscopists varies significantly in different healthcare and resource-level areas of China⁵⁸. The protective effects of colonoscopy and subsequent surveillance strategies may vary according to endoscopist performance. A study assessed 259,885 colonoscopies by 361 endoscopists in the Austria quality assurance program for screening colonoscopies. The study revealed higher CRC mortality rate when the endoscopist had an adenoma detection rate (ADR) < 25%⁵⁹. A current issue in China is a lack of professional endoscopists with high-performance quality, which is especially pronounced in economically underdeveloped areas. The Chinese Preventive Medicine Association released a plan to train physicians and program coordinators from county-level hospitals and cancer screening institutions in 2020 to improve the capacity and performance quality of cancer prevention and management⁶⁰.

Follow-up and endoscopic surveillance

Currently, population-based follow-up for cancer screening in China is inadequate. High-risk individuals' follow-up after baseline examination at most cancer screening sites in China relies on passive linkage with outcomes from the population-based cancer registry or death certification database. Precise follow-up of individuals with abnormal findings after baseline screening is crucial for GI cancer screening with a focus on identifying precursor lesions. Another issue is to release clear recommendations regarding the surveillance intervals. The natural history of esophageal, gastric, and colorectal cancers follows the process from precursor lesions to cancer, which is quite different. Therefore, personalized surveillance intervals should be established for different types of cancer and pathologic characteristics based on the histologic type at baseline screening. There is a lack of high-level evidence in the Chinese population regarding the endoscopic surveillance interval^{28,50,61}.

Screening policies

Population coverage is crucial for the effectiveness of cancer screening in China. Given the large population and limited resources, regular cancer screening of the entire eligible population in China can be unrealistic. Although the National Cancer Screening Program's coverage is gradually expanding, population-level coverage for the top 5 cancers in China is still < 1%^{11,19}. The government should take various measures to increase the proportion of the screening population, including improving the proportion of the awareness of core knowledge on cancer prevention and control among Chinese residents and encouraging opportunistic cancer screening.

Conclusions

Over the past three decades, China has implemented a series of cancer prevention and control measures, contributing to a global reduction in cancer burden. Despite significant progress, continuous exposure to risk factors, such as obesity and population aging in China, may increase the health burden of GI cancers in the future. To address this, China should persist in promoting EC screening in high-incidence areas, explore sensitive biomarkers and risk prediction models for GC and CRC, and investigate non-invasive screening alternatives for endoscopic examination to improve uptake rates. Additionally, efforts should be made to enhance the quality of large-scale population screening data, strengthen data sharing and cooperation, and increase research funding for GI cancers.

Grant support

This work was supported by grants from the Clinical Need Oriented Basic Research Project of Inner Mongolia Academy of Medical Sciences (Grant No. 2023GLLH0132), Scientific Research Fund for the Demonstration Project of Public Hospital Reform and Quality Development (Gastrointestinal Tumor) that is approved by Peking University Cancer Hospital [Inner Mongolia Campus (Grant No. 2023SGGZ068)], and the Beijing Hospitals Authority Youth Programme (Grant No. QML20211102).

Conflict of interest statement

No potential conflicts of interest are disclosed.

Author contributions

Conceived and designed the contents: Xiaotian Zhang and Jiafu Ji.

Retrieved and selected the literature: Xinpu Lu, Fanghui Gu, and Ning Wang.

Wrote the paper: Lei Yang, Li Feng, and Yong Zhu.

References

1. Cancer Today. IARC. [Accessed at 2023 Jul 04]. Available from: <https://gco.iarc.fr/today/home>.
2. Xie Y, Shi L, He X, Luo Y. Gastrointestinal cancers in China, the USA, and Europe. *Gastroenterol Rep (Oxf)*. 2021; 9: 91-104.
3. Liang H, Fan JH, Qiao YL. Epidemiology, etiology, and prevention of esophageal squamous cell carcinoma in China. *Cancer Biol Med*. 2017; 14: 33-41.
4. Lee HW, Huang D, Shin WK, de la Torre K, Song M, Shin A, et al. Frequent low dose alcohol intake increases gastric cancer risk: the Health Examinees-Gem (HEXA-G) study. *Cancer Biol Med*. 2022; 19: 1224-34.
5. Yang L, Ying X, Liu S, Lyu G, Xu Z, Zhang X, et al. Gastric cancer: epidemiology, risk factors and prevention strategies. *Chin J Cancer Res*. 2020; 32: 695-704.
6. Arnold M, Sierra MS, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global patterns and trends in colorectal cancer incidence and mortality. *Gut*. 2017; 66: 683-91.
7. Chen X, Ding J, Li H, Carr PR, Hoffmeister M, Brenner H. The power of a healthy lifestyle for cancer prevention: the example of colorectal cancer. *Cancer Biol Med*. 2022; 19: 1586-97.
8. Qi J, Li M, Wang L, Hu Y, Liu W, Long Z, et al. National and subnational trends in cancer burden in China, 2005-20: an analysis of national mortality surveillance data. *Lancet Public Health*. 2023; 8: e943-55.
9. Edwards BK, Ward E, Kohler BA, Ehemann C, Zauber AG, Anderson RN, et al. Annual report to the nation on the status of cancer, 1975-2006, featuring colorectal cancer trends and impact of interventions (risk factors, screening, and treatment) to reduce future rates. *Cancer*. 2010; 116: 544-73.
10. The Cancer Prevention and Control Outline in China (2004-2010). *China Cancer*. 2004: 65-8.
11. Xia C, Basu P, Kramer BS, Li H, Qu C, Yu XQ, et al. Cancer screening in China: a steep road from evidence to implementation. *Lancet Public Health*. 2023; 8: e996-1005.
12. Plan for Prevention and Control of Chronic Disease in China (2012-15). Bulletin of the Ministry of Health of the People's Republic of China. 2012. [Accessed at 2023 Nov 05]. Available from: https://www.gov.cn/gzdt/2012-05/21/content_2142133.htm.
13. Health China 2030. The state council of the people's republic of China. [Accessed at 2023 Nov 18]. Available from: https://www.gov.cn/zhengce/2016-10/25/content_5124174.htm.
14. Action Plan for Cancer Prevention and Control (2023-30). National Health Commission of the People's Republic of China, 2023. [Accessed at 2023 Dec 01]. Available from: <http://www.nhc.gov.cn/ylyjs/pqt/202311/18bd5bb5abc74ebc896f9d5c9ca63422.shtml>.
15. Zheng R, Zhang S, Zeng H, Wang S, Sun K, Chen R, et al. Cancer incidence and mortality in China, 2016. *J Natl Cancer Cent*. 2022; 2: 1-9.
16. Arnold M, Soerjomataram I, Ferlay J, Forman D. Global incidence of oesophageal cancer by histological subtype in 2012. *Gut*. 2015; 64: 381-7.
17. Cao X, Sun X. Incidence and trend of esophageal cancer. *Chin J Clin Oncol*. 2016; 43: 932-6.
18. Ji J, Li Z, Wang N. Beijing Cancer Registry Annual Report 2022. Beijing, China: Peking University Medical Release; 2023.
19. Zeng H, Chen W, Zheng R, Zhang S, Ji JS, Zou X, et al. Changing cancer survival in China during 2003-15: a pooled analysis of 17 population-based cancer registries. *Lancet Glob Health*. 2018; 6: e555-e67.
20. Chen W, Zeng H, Chen R, Xia R, Yang Z, Xia C, et al. Evaluating efficacy of screening for upper gastrointestinal cancer in China: a study protocol for a randomized controlled trial. *Chin J Cancer Res*. 2017; 29: 294-302.
21. Chen W, Li H, Zheng R, Ren J, Shi J, Cao M, et al. An initial screening strategy based on epidemiologic information in esophageal cancer screening: a prospective evaluation in a community-based cancer screening cohort in rural China. *Gastrointest Endosc*. 2021; 93: 110-8.e2.
22. He Y, Wu Y, Song G, Li Y, Liang D, Jin J, et al. Incidence and mortality rate of esophageal cancer has decreased during past 40 years in Hebei Province, China. *Chin J Cancer Res*. 2015; 27: 562-71.
23. Wei WQ, Chen ZF, He YT, Feng H, Hou J, Lin DM, et al. Long-term follow-up of a community assignment, one-time endoscopic screening study of esophageal cancer in China. *J Clin Oncol*. 2015; 33: 1951-7.
24. Liu M, He Z, Guo C, Xu R, Li F, Ning T, et al. Effectiveness of intensive endoscopic screening for esophageal cancer in China: a community-based study. *Am J Epidemiol*. 2019; 188: 776-84.
25. Yang J, Wei WQ, Niu J, He YT, Liu ZC, Song GH, et al. Estimating the costs of esophageal cancer screening, early diagnosis and treatment in three high risk areas in China. *Asian Pac J Cancer Prev*. 2011; 12: 1245-50.
26. Li F, Liu M, Guo C, Xu R, Li F, Liu Z, et al. Cost-effectiveness of precision screening for esophageal cancer based on individualized risk stratification in China: real-world evidence from the ESECC trial. *Front Oncol*. 2022; 12: 1002693.
27. Wang GQ, Wei WW. [A new transition of the screening, early diagnosis and early treatment project of the upper gastrointestinal

- cancer: opportunistic screening]. *Zhonghua Yu Fang Yi Xue Za Zhi*. 2019; 53: 1084-7.
28. He J, Chen WQ, Li ZS, Li N, Ren JS, Tian JH, et al. China guideline for the screening, early detection and early treatment of esophageal cancer (2022, Beijing). *China Cancer*. 2022; 32: 401-36.
 29. Thrift AP, El-Serag HB. Burden of gastric cancer. *Clin Gastroenterol Hepatol*. 2020; 18: 534-42.
 30. Pan KF, Zhang L, Gerhard M, Ma JL, Liu WD, Ulm K, et al. A large randomised controlled intervention trial to prevent gastric cancer by eradication of *Helicobacter pylori* in Linqu County, China: baseline results and factors affecting the eradication. *Gut*. 2016; 65: 9-18.
 31. Sun X, Mu R, Zhou Y, Dai X, Qiao Y, Zhang S, et al. 1990-1992 mortality of stomach cancer in China. *Zhonghua Zhong Liu Za Zhi*. 2002; 24: 4-8.
 32. Zeng H, Ran X, An L, Zheng R, Zhang S, Ji JS, et al. Disparities in stage at diagnosis for five common cancers in China: a multicentre, hospital-based, observational study. *Lancet Public Health*. 2021; 6: e877-87.
 33. Xu H, Li W. Early detection of gastric cancer in China: progress and opportunities. *Cancer Biol Med*. 2022; 19: 1622-8.
 34. Fan X, Qin X, Zhang Y, Li Z, Zhou T, Zhang J, et al. Screening for gastric cancer in China: advances, challenges and visions. *Chin J Cancer Res*. 2021; 33: 168-80.
 35. Chen R, Liu Y, Song G, Li B, Zhao D, Hua Z, et al. Effectiveness of one-time endoscopic screening programme in prevention of upper gastrointestinal cancer in China: a multicentre population-based cohort study. *Gut*. 2021; 70: 251-60.
 36. Li WQ, Qin XX, Li ZX, Wang LH, Liu ZC, Fan XH, et al. Beneficial effects of endoscopic screening on gastric cancer and optimal screening interval: a population-based study. *Endoscopy*. 2022; 54: 848-58.
 37. Leja M, You W, Camargo MC, Saito H. Implementation of gastric cancer screening - the global experience. *Best Pract Res Clin Gastroenterol*. 2014; 28: 1093-106.
 38. Cai Q, Zhu C, Yuan Y, Feng Q, Feng Y, Hao Y, et al. Development and validation of a prediction rule for estimating gastric cancer risk in the Chinese high-risk population: a nationwide multicentre study. *Gut*. 2019; 68: 1576-87.
 39. Ford AC, Yuan Y, Moayyedi P. *Helicobacter pylori* eradication therapy to prevent gastric cancer: systematic review and meta-analysis. *Gut*. 2020; 69: 2113-21.
 40. Joseph DA, Meester RG, Zauber AG, Manninen DL, Wings L, Dong FB, et al. Colorectal cancer screening: estimated future colonoscopy need and current volume and capacity. *Cancer*. 2016; 122: 2479-86.
 41. Meester RG, Doubeni CA, Zauber AG, Goede SL, Levin TR, Corley DA, et al. Public health impact of achieving 80% colorectal cancer screening rates in the United States by 2018. *Cancer*. 2015; 121: 2281-5.
 42. Siegel RL, Miller KD, Wagle NS, Jemal A. Cancer statistics, 2023. *CA Cancer J Clin*. 2023; 73: 17-48.
 43. Carethers JM, Jung BH. Genetics and genetic biomarkers in sporadic colorectal cancer. *Gastroenterology*. 2015; 149: 1177-90 e3.
 44. Kuipers EJ, Rösch T, Bretthauer M. Colorectal cancer screening - optimizing current strategies and new directions. *Nat Rev Clin Oncol*. 2013; 10: 130-42.
 45. Zheng S, Chen K, Liu X, Ma X, Yu H, Chen K, et al. Cluster randomization trial of sequence mass screening for colorectal cancer. *Dis Colon Rectum*. 2003; 46: 51-8.
 46. Dai M, Shi J, Li N. The design and objective for Cancer Screening Program in Urban China. *Chin J Prev Med*. 2013; 47: 179-82.
 47. Chen H, Lu B, Dai M. Colorectal cancer screening in China: status, challenges, and prospects - China, 2022. *China CDC Wkly*. 2022; 4: 322-8.
 48. Li N, Zhou YY, Lu M, Zhang YH, Lu B, Luo CY, et al. Participation rate and detection of colorectal neoplasms based on multi-round fecal immunochemical testing for colorectal cancer screening in the Chinese population. *Zhonghua Zhong Liu Za Zhi*. 2023; 45: 1041-50.
 49. Imperiale TF, Ransohoff DF, Itzkowitz SH, Levin TR, Lavin P, Lidgard GP, et al. Multitarget stool DNA testing for colorectal-cancer screening. *N Engl J Med*. 2014; 370: 1287-97.
 50. National Cancer Center, China, Expert Group of the Development of China Guideline for the Screening, Early Detection and Early Treatment of Colorectal Cancer. China guideline for the screening, early detection and early treatment of colorectal cancer (2020, Beijing). *Zhonghua Zhong Liu Za Zhi*. 2021; 43: 16-38.
 51. Chen H, Lu M, Liu C, Zou S, Du L, Liao X, et al. Comparative evaluation of participation and diagnostic yield of colonoscopy vs fecal immunochemical test vs risk-adapted screening in colorectal cancer screening: interim analysis of a multicenter randomized controlled trial (TARGET-C). *Am J Gastroenterol*. 2020; 115: 1264-74.
 52. Luo H, Xu G, Li C, He L, Luo L, Wang Z, et al. Real-time artificial intelligence for detection of upper gastrointestinal cancer by endoscopy: a multicentre, case-control, diagnostic study. *Lancet Oncol*. 2019; 20: 1645-54.
 53. Lei Y, Xi Z, Shuo L, Huichao L. Gastric cancer screening of early diagnosis and treatment of cancer project in urban Beijing from 2014 to 2019. *Chin J Cancer Prev Treat*. 2023; 30: 1414-9.
 54. Fanghui G, Lei Y, Xi Z, Qingyu L. Analysis on the complacance with endoscopic screening for colorectal cancer among high-risk population in Beijing. *Chin Prev Med*. 2021; 22: 428-33.
 55. Xi Z, Lei Y, Qingyu L, Shuo L. Analysis of screening results and compliance of colorectal cancer among rural residents in Beijing, 2017-2021. *China Cancer*. 2023; 32: 603-9.
 56. Zhang X, Yang L, Liu S, Li H, Li Q, Li H, et al. Performance of different colorectal cancer screening strategies: a long-term passive follow-up population-based screening program in Beijing, China. *BMC Public Health*. 2023; 23: 1640.
 57. Bretthauer M, Løberg M, Wieszczy P, Kalager M, Emilsson L, Garborg K, et al. Effect of colonoscopy screening on risks of

- colorectal cancer and related death. *N Engl J Med.* 2022; 387: 1547-56.
58. Jibin L, Wenqiang W, Liwei Z, Yuqin L. Skill assessment results and influencing factors among endoscopists of upper gastrointestinal cancer screening program in rural China. *China Cancer.* 2021; 30: 445-51.
59. Waldmann E, Kammerlander AA, Gessl I, Penz D, Majcher B, Hinterberger A, et al. Association of adenoma detection rate and adenoma characteristics with colorectal cancer mortality after screening colonoscopy. *Clin Gastroenterol Hepatol.* 2021; 19: 1890-8.
60. Non-communicable Disease Management - Training Program for Cancer Screening and Early Diagnosis 2020. [Accessed at 2023 Nov 11]. Available from: <https://www.cpma.org.cn/zhyfyx/xhdt/202008/5cf9c773e48346338550ea75baa7bc41.shtml>.
61. Jie H, Wanqing C, Zhaoshen L, Ni L. China guideline for the screening, early detection and early treatment of gastric cancer (2022, Beijing). *China Cancer.* 2022; 31: 488-527.

Cite this article as: Yang L, Feng L, Zhu Y, Wang N, Lu X, Gu F, et al. Reducing the global cancer burden with gastrointestinal screening: China's 30 years practice. *Cancer Biol Med.* 2024; 21: 204-212. doi: 10.20892/j.issn.2095-3941.2023.0516