



## ORIGINAL ARTICLE

# DNA damage response-related immune activation signature predicts the response to immune checkpoint inhibitors: from gastrointestinal cancer analysis to pan-cancer validation

Junya Yan<sup>1,2\*</sup>, Shibo Wang<sup>1\*</sup>, Jing Zhang<sup>3\*</sup>, Qiangqiang Yuan<sup>1</sup>, Xianchun Gao<sup>1</sup>, Nannan Zhang<sup>1</sup>, Yan Pan<sup>1</sup>, Haohao Zhang<sup>1</sup>, Kun Liu<sup>4</sup>, Jun Yu<sup>1</sup>, Linbin Lu<sup>1</sup>, Hui Liu<sup>5</sup>, Xiaoliang Gao<sup>1</sup>, Sheng Zhao<sup>1</sup>, Wenyao Zhang<sup>1</sup>, Abudurousuli Reyila<sup>1</sup>, Yu Qi<sup>1</sup>, Qiuji Zhang<sup>6</sup>, Shundong Cang<sup>2</sup>, Yuanyuan Lu<sup>1</sup>, Yanglin Pan<sup>1</sup>, Yan Kong<sup>7</sup>, Yongzhan Nie<sup>1</sup>

<sup>1</sup>State Key Laboratory of Holistic Integrative Management of Gastrointestinal Cancers and National Clinical Research Center for Digestive Diseases, Xijing Hospital of Air Force Military Medical University, Xi'an 710032, China; <sup>2</sup>Department of Oncology, Henan Provincial People's Hospital, Zhengzhou University People's Hospital, Zhengzhou 450003, China; <sup>3</sup>Faculty of Life Science, Northwest University, Xi'an 710069, China; <sup>4</sup>Unit 73211 of the People's Liberation Army, Nanjing 211800, China; <sup>5</sup>Key Laboratory of Resource Biology and Biotechnology in Western China (Ministry of Education), School of Medicine, Northwest University, Xi'an 710069, China; <sup>6</sup>Shaanxi University of Chinese Medicine, Second Clinical Medicine Faculty, Xi'an 712046, China; <sup>7</sup>Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education), Department of Renal Cancer and Melanoma, Peking University Cancer Hospital and Institute, Beijing 100142, China

**ABSTRACT**

**Objective:** DNA damage response (DDR) deficiency has emerged as a prominent determinant of tumor immunogenicity. This study aimed to construct a DDR-related immune activation (DRIA) signature and evaluate the predictive accuracy of the DRIA signature for response to immune checkpoint inhibitor (ICI) therapy in gastrointestinal (GI) cancer.

**Methods:** A DRIA signature was established based on two previously reported DNA damage immune response assays. Clinical and gene expression data from two published GI cancer cohorts were used to assess and validate the association between the DRIA score and response to ICI therapy. The predictive accuracy of the DRIA score was validated based on one ICI-treated melanoma and three pan-cancer published cohorts.

**Results:** The DRIA signature includes three genes (*CXCL10*, *IDO1*, and *IFI44L*). In the discovery cancer cohort, DRIA-high patients with gastric cancer achieved a higher response rate to ICI therapy than DRIA-low patients (81.8% vs. 8.8%;  $P < 0.001$ ), and the predictive accuracy of the DRIA score [area under the receiver operating characteristic curve (AUC) = 0.845] was superior to the predictive accuracy of PD-L1 expression, tumor mutational burden, microsatellite instability, and Epstein-Barr virus status. The validation cohort demonstrated that the DRIA score identified responders with microsatellite-stable colorectal and pancreatic adenocarcinoma who received dual PD-1 and CTLA-4 blockade with radiation therapy. Furthermore, the predictive performance of the DRIA score was shown to be robust through an extended validation in melanoma, urothelial cancer, and pan-cancer.

**Conclusions:** The DRIA signature has superior and robust predictive accuracy for the efficacy of ICI therapy in GI cancer and pan-cancer, indicating that the DRIA signature may serve as a powerful biomarker for guiding ICI therapy decisions.

**KEYWORDS**

DNA damage response-related immune activation; immune checkpoint inhibitors; biomarker; gastrointestinal cancer; pan-cancer

\*These authors contributed equally to this work.

Correspondence to: Yan Kong and Yongzhan Nie  
E-mail: k-yan08@163.com and yongznjie@fmmu.edu.cn  
ORCID ID: <https://orcid.org/0000-0002-6368-2112>  
and <https://orcid.org/0000-0003-1124-4548>

Received August 15, 2023; accepted November 13, 2023;  
published online December 29, 2023.

Available at [www.cancerbiomed.org](http://www.cancerbiomed.org)

©2024 Cancer Biology & Medicine. Creative Commons  
Attribution-NonCommercial 4.0 International License

## Introduction

Gastrointestinal (GI) cancer has the highest disease burden among all malignancies with global incidence and mortality rates of 26% and 35%, respectively<sup>1</sup>. Traditional treatments for GI cancer, including surgical resection, chemotherapy, radiotherapy, and targeted therapy, have not achieved satisfactory outcomes<sup>2</sup>. In recent years the development of immunotherapy has revolutionized the therapeutic landscape

for multiple cancer types, including a subset of GI cancers<sup>3</sup>. Immune checkpoint inhibitor (ICI) targeting PD-1/PD-L1 signaling have been approved as standard first-line therapy for advanced esophagogastric and colorectal cancer<sup>4,5</sup>. The objective response rate (ORR) of ICI therapy, however, is only 10%–20% in patients with GI cancer<sup>6</sup>, highlighting the urgent need to identify optimal biomarkers for precise treatment.

Notably, some biomarkers have been developed to predict the efficacy of ICI therapy in patients with GI cancer, including PD-L1 expression<sup>7</sup>, tumor mutational burden (TMB)<sup>8</sup>, microsatellite instability (MSI) status<sup>9</sup>, and Epstein–Barr virus (EBV) status<sup>10</sup>. To date, only MSI-high (MSI-H) has been confirmed to be a useful predictive biomarker for GI cancer. Nevertheless, the MSI-H subtype is present in only a small proportion (0%–5%) of colorectal cancers<sup>11</sup>. PD-L1 expression is one of the most widely studied immunotherapy biomarkers; however, clinical trials have yielded controversial results<sup>7,12,13</sup>, which limits the predictive value of PD-L1 expression in patients with GI cancer. In addition, TMB, as a biomarker of GI cancer, faces several challenges, such as platform uniformity, intra-tumoral heterogeneity, and lack of consensus on thresholds<sup>6,14,15</sup>. Overall, the currently available biomarkers do not meet the criteria for clinical application in ICI therapy for GI cancer.

Genomic instability, as one of the hallmarks of various cancers, is mainly caused by the increasing accumulation of damaged DNA and DNA damage response (DDR) deficiency<sup>16</sup>. Although DDR deficiency drives genomic instability and tumor progression, DDR deficiency also provides potential therapeutic opportunities. The excellent efficacy of olaparib in ovarian cancer patients with *BRCA1/2* mutations is probably the best example of this interplay<sup>17</sup>. In addition, data from multiple studies have shown that patients responding to ICI therapy commonly harbor *BRCA2* mutations<sup>18</sup>. Other investigations also demonstrated that deleterious alterations in genes involved in DDR pathways induce a hyper-mutational phenotype and improve the survival outcomes following ICI therapy<sup>19,20</sup>. DDR impairment can be induced not only by genomic alterations, but also epigenetic changes in DDR pathways<sup>21</sup>. Indeed, both mechanisms underlying DDR impairment induce the accumulation of DNA damage and activate the cyclic GMP-AMP synthase (cGAS)-stimulator of interferon genes (STING) pathway to create an inflammatory microenvironment<sup>21</sup>, thereby defining a distinct subgroup of patients suitable for ICI therapy.

Notably, a recently developed immune-driven 44-gene signature [DNA damage immune response (DDIR) assay<sup>22</sup>],

summarized the common immune active process from the accumulation of DNA damage, which created an inflammatory microenvironment characterized by increased CD8<sup>+</sup> T cell infiltration and PD-L1 expression<sup>23</sup>. Furthermore, DDIR positivity has been reported to predict the clinical response to chemotherapy among patients with breast cancer<sup>24,25</sup> and esophageal adenocarcinoma<sup>26</sup>. Although the DDIR signature cannot predict an improved response to oxaliplatin chemotherapy in patients with colorectal cancer, a refined 9-gene DDIR signature showed a strong association with MSI and the consensus molecular subtype<sup>27</sup>. Based on the above analyses, we propose the DDIR signature potential as a classifier in identifying GI cancers that may benefit from ICI therapy.

In the current study we first established a novel DDR-related immune activation (DRIA) signature by detecting a smaller panel of genes that can be easily translated into an easy-to-use clinical assay. By collecting the clinical and transcriptome profile data of two ICI-treated GI cancer cohorts, we explored and validated the association between the DRIA signature and clinical response to ICI therapy. To broaden the applicable certificate of DRIA, we retrospectively collected one ICI-treated melanoma cohort from our center and three pan-cancer cohorts from published datasets for extended validation of the DRIA predictive robustness. This study developed and revealed a novel signature (DRIA) consisting of three genes that better predict therapeutic efficacy of ICI therapy in GI cancer and pan-cancer.

## Materials and methods

### Patients and transcriptome profile data

Fifty-five patients with melanoma who received anti-PD-1 monotherapy between March 2016 and March 2019 were recruited for this study from Peking University Cancer Hospital (PUCH). Formalin-fixed paraffin-embedded (FFPE) biopsy specimens were collected from each patient before undergoing immunotherapy. Whole-transcriptome RNA sequencing was performed on the Illumina NovaSeq 6000 platform (San Diego, California, USA). The details of processing the gene expression data are described in our previous study<sup>28</sup>.

All the clinical and pathologic data, including age, gender, primary site, metastasis, and clinical response, were extracted from the medical record review. Patient response was determined using the Response Evaluation Criteria in Solid Tumors [RECIST (version 1.1)] criteria as follows: complete response

(CR); partial response (PR); stable disease (SD); and progressive disease (PD). The ORR was defined as the proportion of patients with a CR or PR. Progression-free survival (PFS) was defined as the time from the initiation of ICI therapy to the date of disease progression or death from any cause. Patients without disease progression were censored on the date of their last scan. Overall survival (OS) was calculated from the date ICI therapy commenced to the date of death. Patients who did not die were censored on the date of their last scan. In this study, the ORR, PFS, and OS were the primary clinical outcomes. This study was approved by the Institutional Review Board of PUCH (2019KT92) and was in accordance with the Declaration of Helsinki. Written informed consent was obtained from all participants.

### External cohort acquisition

The RNA-seq and clinical data from five publicly available cohorts treated with ICI therapy were collected for analysis in this study, as follows: (1) the Kim18 cohort consisted of 45 patients with metastatic gastric cancer who received anti-PD-1 therapy at the Samsung Medical Center<sup>10</sup>; (2) the Parikh22 cohort included 22 patients with metastatic microsatellite-stable (MSS) colorectal and pancreatic adenocarcinoma who received dual PD-1 and CTLA-4 blockade with radiation therapy in a phase II trial (NCT03104439)<sup>29</sup>; (3) the Gide19 cohort was comprised of 41 patients with advanced melanoma treated with anti-PD-1 therapy at the University of Sydney<sup>30</sup>; (4) the IMvigor210 cohort included 298 patients with metastatic urothelial cancer who received anti-PD-L1 therapy at the Memorial Sloan Kettering Cancer Center<sup>31</sup>; and (5) the Pender21 cohort comprised 56 pan-cancer patients across 17 solid tumor types treated with anti-PD-(L)1 therapy at the University of British Columbia<sup>32</sup>. Detailed information for the five immunotherapy cohorts is summarized in **Table S1**.

### Definition of the DRIA signature

Two previously reported DDIR signatures in breast and colorectal cancers<sup>22,27</sup> contain 44 genes and 9 genes, respectively (**Table S2**). Three overlapping genes between the two DDIR signatures were defined as the DRIA signature, and the DRIA score was calculated as the geometric mean of the three genes. Based on the DRIA scores and immunotherapy responses, receiver operating characteristic (ROC) curve analysis was performed and the cut-off value that generated the maximum Youden index was used to define

DRIA status. A DRIA score greater than the threshold was classified as DRIA-high and a DRIA score less than the threshold was classified as DRIA-low.

### The Cancer Genome Atlas (TCGA) cohort

We downloaded the transcriptome profiling and survival data of patients with GI cancer (esophageal cancer,  $n = 161$ ; gastric cancer,  $n = 375$ ; and colorectal cancer,  $n = 622$ ) from the TCGA database (<http://xena.ucsc.edu/>). Some immune-related molecular features, including the PD-L1 score, TMB, MSI status, CD8A score, and IFN- $\gamma$  score were extracted from cBioPortal (<https://www.cbioportal.org/>) and the TCGA-GDC website (<https://gdc.cancer.gov/about-data/publications/panimmune>) to analyze the correlations with the DRIA signature.

### Assessment of DDR mutation status

Previous studies defined a DDR gene list (**Table S3**), including 34 genes involved in 6 major DDR pathways<sup>19,20,33</sup>. According to the co-mutation counts of these DDR genes, the patients were divided into DDR-deficient ( $\geq 2$  mutations) and DDR-proficient groups ( $< 2$  mutations). The mutation information regarding the 34 genes in TCGA-STAD cohort were retrieved from the cBioPortal for Cancer Genomics (<https://www.cbioportal.org/>). The non-synonymous mutations included TRUNC (Frameshift del, Frameshift ins, nonsense, nonstop, splice region, splice site), INFRAME (Inframe del and Inframe ins), and MISSENSE mutations.

### Biomarker analysis

Detailed information regarding clinically known biomarkers for immunotherapy were available in the Kim18 and IMvigor210 cohorts<sup>10,31</sup>. PD-L1 expression [combined positive score (CPS)], MSI status, TMB, and EBV status data were extracted from the Kim18 cohort. Data from the IMvigor210 cohort with respect to CD8<sup>+</sup> T cell infiltration, PD-L1 expression in tumor-infiltrating immune cells (ICs) and tumor cells (TCs), TMB, and tumor neoantigen burden (TNB) were also separated. We performed correlation analyses between these biomarkers and the DRIA score, and compared the predictive accuracy to determine the efficacy of immunotherapy. We also included published signatures (**Table S4**) based on the transcriptome profile data and evaluated the predictive performance based on the Kim18 cohort response to immunotherapy.

## Statistical analyses

SPSS (version 23.0) and GraphPad Prism software (version 8) were used for statistical analyses and graphical representations. For normally distributed continuous variables, a t-test was used to compare mean values, whereas the Mann–Whitney U test was used for non-normally distributed continuous variables. Correlations between the DRIA status and categorical measurements, such as tumor response, were assessed using a chi-square test or Fisher's exact test. Kaplan–Meier estimates of survival outcomes, including PFS and OS, were calculated using a log-rank test. ROC curve analyses were performed to evaluate the predictive accuracy of the DRIA score and other biomarkers with respect to ICI therapy. For all statistical tests, a  $P < 0.05$  (two-tailed test) was considered statistically significant.

## Results

### Immune characteristics and prognosis of DRIA in GI cancer

We identified three overlapping genes (*CXCL10*, *IDO1*, and *IFI44L*) in the two previously reported DDIR signatures (Table S2) and defined the overlapping genes as the DRIA signature. Using these three consensus genes to generate an unweighted cumulative DRIA score, we observed strong positive correlations between the DRIA score and the two original DDIR scores in the TCGA-GI cancer cohort (44-gene: Spearman  $r = 0.621$ , Figure S1A; 9-gene: Spearman  $r = 0.963$ , Figure S1B). Notably, the level of *CXCL10*, *IDO1*, and *IFI44L* gene expression was increased in DDR-deficient tumors compared with DDR-proficient tumors (Figure S2A–C), which indicated the necessity of the three genes for DDIR characteristics. By analyzing the immune-related molecular features of the TCGA-GI cancer cohort, we demonstrated a markedly increased DRIA score in MSI patients compared to MSS patients (Figure S1C). The DRIA score was positively correlated with the TMB (Spearman  $r = 0.154$ , Figure S1D), PD-L1 score (Spearman  $r = 0.695$ , Figure S1E), CD8A score (Spearman  $r = 0.737$ , Figure S1F), and IFN- $\gamma$  score (Spearman  $r = 0.832$ , Figure S1G). Therefore, patients with different DRIA scores had distinct tumor immune microenvironment characteristics, indicating that the DRIA signature has the potential to predict the response to ICI therapy in patients with GI cancer.

Survival analyses were performed according to DRIA status in the TCGA-GI cancer cohort to investigate the potential

prognostic role of DRIA. No significant survival difference was observed between the DRIA-high and -low subsets in patients with esophageal, gastric, and colorectal cancers without ICI therapy (Figure S3A–C), indicating that DRIA was not a prognostic factor.


### Exploration of the association between DRIA and response to ICI therapy in gastric cancer

The flow diagram depicting the multi-cohort analysis of the DRIA predictive value for the efficacy of ICI therapy is presented in Figure 1. A total of 517 patients treated with ICI from 6 cohorts, including GI cancer, melanoma, urothelial cancer, and pan-cancer, were included in this study. The clinical baseline characteristics are summarized in Table S5. The Kim18 cohort ( $n = 45$ ) served as the discovery cohort with which to determine the association between DRIA and the efficacy of ICI therapy in patients with GI cancer. Furthermore, we used the Parikh22 cohort ( $n = 22$ ) to validate the predictive value of DRIA in patients with GI cancer. Finally, pan-cancer extended validations of the DRIA were performed in the PUCH ( $n = 55$ ), Gide19 ( $n = 41$ ), IMvigor210 ( $n = 298$ ), and Pender21 ( $n = 56$ ) cohorts.

The median DRIA score of the patients in the CR/PR group was significantly higher than the patients in the SD/PD group of the Kim18 cohort (Figure 2A). Next, we integrated the DRIA score with the clinical response data to create an ROC curve and quantify the prediction accuracy. The results revealed that the area under the ROC curve (AUC) was 0.838 (95% CI, 0.680–0.997; Figure 2B). The patients were divided into a DRIA-high group ( $n = 11$ ) and a DRIA-low group ( $n = 34$ ) using a cut-off value that generated the maximum Youden index. The DRIA-high group had a markedly higher ORR than the DRIA-low group (81.8% vs. 8.8%; Figure 2C). Moreover, the waterfall plot demonstrated a positive correlation between the DRIA score and a favorable clinical response (Figure 2D). These results supported DRIA as a superior predictor of the clinical benefits of ICI therapy in patients with gastric cancer.

### Comparison of DRIA with known clinical biomarkers in gastric cancer

Several emerging predictors have been developed for gastric cancer immunotherapy, including PD-L1 expression, MSI status, EBV status, and TMB. We performed correlation analyses between DRIA status and these biomarkers based




**Figure 1** Flow diagram of this study. Flow diagram shows the analytic process of the DRIA predictive value in multiple ICI-treated cohorts. DRIA, DNA damage response-related immune activation; ICIs, immune checkpoint inhibitors; GI, gastrointestinal; MSS, microsatellite-stable.

on the available data in the Kim18 cohort. Detailed information regarding the DRIA status and other biomarkers in each patient is shown in a heatmap (**Figure 3A**). When patients were classified into 3 groups according to the PD-L1 CPS, the  $\text{CPS} \geq 10$  group had a significantly increased DRIA score combined with the  $1 \leq \text{CPS} < 10$  and  $\text{CPS} = 0$  groups (**Figure 3B**). Correlation analysis also showed a remarkably positive correlation between the PD-L1 CPS and the DRIA score (Spearman  $r = 0.385$ ; **Figure 3C**). Additionally, the median DRIA score was significantly higher in EBV-positive patients than EBV-negative patients (**Figure 3D**). Similarly, MSI-H and TMB-high patients had relatively increased DRIA scores (**Figure 3E, F**), although the difference did not reach significance. Furthermore, we applied ROC curve analysis to compare the DRIA predictive accuracy and these biomarkers. The DRIA predicting AUC was 0.845, which was higher than PD-L1, TMB, MSI, and EBV (**Figure 3G**). We further determined whether synergizing DRIA with these biomarkers enhanced the prediction accuracy of the

response to ICI therapy. The pairwise combination of DRIA with PD-L1, TMB, MSI, and EBV showed improved patient stratification (**Figure S4A**). Remarkably, the predicting AUC of the DRIA with PD-L1 combination reached 0.947 (95% CI, 0.876–1.000). According to the scoring system constructed by incorporating DRIA and PD-L1, patients were stratified into 3 groups (score = 0, 1, and 2). The proportion of patients who achieved a CR/PR was 90.0%, 15.4%, and 0% for the 2, 1, and 0 score groups, respectively (**Figure S4B**).

Additionally, the expression signature of the gene series has been used to predict the response to ICI therapy. Therefore, we investigated the association between the DRIA score and these signature scores in the Kim18 cohort. The heatmap showed that these signature scores were dramatically upregulated in the DRIA-high group compared to the DRIA-low group (**Figure 3H**). We also performed ROC curve analyses. The DRIA predicting AUC was higher than these signatures, indicating that the DRIA signature had a stronger predictive ability



**Figure 2** Association between DRIA and the clinical response to ICI therapy in gastric cancer. (A) Distribution of DRIA scores for distinct clinical response groups in the Kim18-gastric cancer cohort. (B) ROC curve of sensitivity versus 1-specificity for the DRIA score predictive value to the clinical response. (C) Stacked bar plot depicting different fractions of clinical response for patients in DRIA-high and -low groups. (D) Waterfall plot of DRIA score for distinct clinical response groups. DRIA, DNA damage response-related immune activation; ICIs, immune checkpoint inhibitors; CR, complete response; PR, partial response; SD, stable disease; PD, progressive disease; AUC, area under the ROC curve; CI, confidence interval. \*\*\* $P < 0.001$ .

for ICI therapy than these previously published immune-related signatures (Figure 3I).

### Validation of the DRIA predictive performance in MSS colorectal cancer and pancreatic adenocarcinoma

The addition of radiation therapy to dual PD-1 and CTLA-4 blockade has demonstrated some efficacy in MSS colorectal cancer and pancreatic adenocarcinoma<sup>29</sup>. Therefore, we attempted to validate the DRIA predictive value for the clinical benefits of this combination therapy in the Parikh22 cohort. ROC curve analysis indicated that the predictive accuracy was excellent, with an AUC of 0.806 (95% CI, 0.580–1.000; Figure 4A). The ORR in the DRIA-high group was superior

to the DRIA-low group (42.9% vs. 0%; Figure 4B). Notably, the waterfall plot suggested that all patients responsive to combination therapy were in the DRIA-high group (Figure 4C). As expected, the DRIA-low patients had a worse PFS (median PFS, 1.77 vs. 2.76 months; Figure 4D) and OS (median OS, 6.14 vs. 11.73 months; Figure 4E) than DRIA-high patients.

### Extended validation of the DRIA predictive robustness for ICI therapy efficacy in pan-cancer cohorts

To confirm the robustness of DRIA in predicting the benefits of ICI therapy, we included four immunotherapy cohorts with multiple types of cancer (melanoma, urothelial cancer, and pan-cancer) for subsequent analyses. Patients in the

CR/PR group tended to have higher DRIA scores than the PD/SD group in the PUCH, which is consistent with the two GI cancer (Figure 5A), Gide19 (Figure 5B), Pender19 (Figure 5C), and IMvigor210 (Figure S5A) immunotherapy cohorts. The waterfall plot showed a positive association between a higher DRIA score and a better clinical response (Figure S6A–C). ROC curve analyses revealed that the resulting AUCs were 0.760 (95% CI, 0.612–0.907; Figure 5D), 0.845 (95% CI, 0.715–0.974; Figure 5E), and 0.766 (95% CI, 0.596–0.936; Figure 5F), respectively. In addition, patients in the DRIA-high group had a better ORR than the DRIA-low group in the PUCH (40.0% vs. 8.0%; Figure 5G), Gide19 (78.9% vs. 18.2%; Figure 5H), and Pender21 (50.0% vs. 6.8%; Figure 5I) cohorts. Furthermore, we compared the clinical outcomes between the two DRIA groups in the four cohorts. The DRIA-low group had a significantly poorer PFS (Figure 6A–C) and OS (Figures S5B and S7A, B) than the DRIA-high group. In agreement with these results, the multivariate Cox regression analyses showed that the higher DRIA score was an independent prognostic factor for predicting a favorable PFS (Figure 6D–F) and OS (Figures S5C and S7C, D) in patients receiving ICI therapy.

We also validated the correlation between the DRIA and these biomarkers based on the available CD8<sup>+</sup> T cell infiltration, PD-L1 expression, TMB, and TNB data in the IMvigor210 cohort. When the tumors were divided into three classic immune phenotypes according to CD8<sup>+</sup> T cells infiltration, we observed that patients with an immune-inflamed phenotype had the highest DRIA scores compared to the other phenotypes (Figure S5D). As predicted, the DRIA-high patients had a relatively higher TMB (Figure S5E) and TNB (Figure S5F). Similarly, a higher DRIA score was positively correlated with higher PD-L1 expression (Figure S5G, H). Above all, the results of the four immunotherapy cohorts fully demonstrated that DRIA precisely predicts the clinical benefit of ICI therapy in pan-cancer.

### Comparison of the DRIA predictive performance with two original DDIR signatures

To facilitate the clinical translation of DRIA, we further conducted ROC analyses to compare the predictive performance of DRIA with two original DDIR signatures as a function of






Figure 3 Continued



**Figure 3** Comparison of the predictive accuracy of DRIA with known clinical biomarkers in gastric cancer. (A) Heatmap of DRIA distributions and known clinical biomarkers for distinct clinical response groups in the Kim18-gastric cancer cohort. (B) Comparison of the DRIA score in three groups according to the level of PD-L1 expression. (C) Correlation of the DRIA score with the level of PD-L1 expression. (D-F) Comparison of the DRIA score in different groups according to EBV (D), MSI (E), and TMB status (F). (G) ROC curves measuring the predictive accuracy of DRIA, PD-L1, TMB, MSI, and EBV for response to ICI therapy. (H) Heatmap showing differently expressed immune-related signatures between the DRIA-high and -low groups. (I) Comparison of the DRIA AUCs and seven published immune-related signatures in predicting the response to ICI therapy. DRIA, DNA damage response-related immune activation; BOR, best of response; MSI, microsatellite instability; EBV, Epstein-Barr virus; TMB, tumor mutational burden; CR, complete response; PR, partial response; SD, stable disease; PD, progressive disease; CPS, combined positive score; ICIs, immune checkpoint inhibitors; AUC, area under the ROC curve; TLS, tertiary lymphoid structures; NA, not available. \* $P < 0.05$ , \*\* $P < 0.01$ .

response to ICI therapy in five immunotherapy cohorts. The predicting DRIA AUCs were essentially equivalent to the 9- and 44-gene DDIR signatures (Table S6). Taken together, the

DRIA signature conferred superior predictive accuracy for ICI therapy efficacy, which warrants further investigations in prospective clinical trials.




**Figure 4** Correlation of DRIA to clinical benefit from ICI-based combination therapy in patients with MSS GI cancer. (A) ROC curve of sensitivity versus 1-specificity for the DRIA score predictive value to clinical response in the Parikh22-MSS GI cancer cohort. (B) Comparison of objective response rate of patients according to the DRIA status. (C) The DRIA scores of individual patients and their clinical responses to ICI therapy. (D-E) Kaplan–Meier survival curves comparing PFS (D) and OS (E) between DRIA-high and -low groups. DRIA, DNA damage response-related immune activation; ICIs, immune checkpoint inhibitors; MSS, microsatellite-stable; GI, gastrointestinal; AUC, area under the ROC curve; CI, confidence interval; CR, complete response; PR, partial response; SD, stable disease; PD, progressive disease; NE, not evaluated; PFS, progression-free survival; OS, overall survival.

## Discussion

In this study we developed and validated a novel immune gene signature, the DRIA, which consists of three genes that robustly predict clinical benefit from ICI therapy in GI cancer and pan-cancer patients. Our results revealed that DRIA positivity effectively identifies patients with improved response rates and survival outcomes in the context of ICI therapy. Indeed, the predictive accuracy of DRIA was superior to the known clinical biomarkers in GI cancer, such as PD-L1 and TMB.

Currently, development of effective biomarkers is required to guide ICI therapy decisions for different types of cancer, including GI cancer. A series of biomarkers, including PD-L1,

TMB, MSI, and EBV, have been widely used in the clinical management of patients with GI cancer<sup>6</sup>. These biomarkers have encountered several issues<sup>6,7,11-13</sup>, such as prediction instability, intra-tumoral heterogeneity, and fewer predicted benefits to populations, which incites us to explore more markers with universal mechanisms. DDR deficiency has recently emerged to increase the level of neoantigens and tumor immunogenicity, and activate the cGAS-STING pathway to reshape the tumor microenvironment, ultimately leading to the effectiveness of ICI therapy<sup>21</sup>. In view of the high frequency of DDR deficiency in tumorigenesis and progression, extensive efforts have been made to identify DDR-related biomarkers for predicting the response to ICI therapy. Several investigations have demonstrated that gene mutations



**Figure 5** Validation of the predictive robustness of DRIA in ICI-treated pan-cancer cohorts. (A-C) Comparison of the DRIA scores between responders and non-responders in PUCH melanoma (A), Gide19 melanoma (B), and Pender21 pan-cancer cohorts (C). (D-F) ROC curves measuring the predictive value of the DRIA score in PUCH melanoma (D), Gide19 melanoma (E), and Pender21 pan-cancer cohorts (F). (G-I) Proportion of objective response for patients with DRIA-high and -low status in PUCH melanoma (G), Gide19 melanoma (H), and Pender21 pan-cancer cohorts (I). DRIA, DNA damage response-related immune activation; ICIs, immune checkpoint inhibitors; CR, complete response; PR, partial response; SD, stable disease; PD, progressive disease; AUC, area under the ROC curve; CI, confidence interval. \* $P < 0.05$ ; \*\* $P < 0.01$ ; \*\*\* $P < 0.001$ .

in different DDR pathways result in a durable clinical benefit from ICI therapy in various types of cancer, including melanoma<sup>34</sup>, lung cancer<sup>35</sup>, bladder cancer<sup>19</sup>, and GI cancer<sup>20</sup>. Previous studies, however, have also shown that even though *BRCA1/2* mutations may confer sensitivity to ICI therapy, this is not held in common for all patients because not all mutations induce DDR deficiency or compensate for by alternate mechanisms<sup>36</sup>. Conversely, *BRCA1/2* wide-type tumors confers a DDR-deficient phenotype due to epigenetic silencing of *BRCA1/2*<sup>37,38</sup>. Therefore, a transcriptome-based signature which assesses the downstream effects of genomic and epigenetic changes of altered DDR pathways may more accurately predict the sensitivity to ICI therapy.

The DDIR signature summarizes the cascade of changes in immune activation from the resulting accumulation of DNA damage<sup>22,23</sup>. Therefore, the DDIR signature could serve as a promising marker to predict the clinical response to ICI therapy. Considering the relatively larger panel of genes in the

original DDIR signature that cannot be easily translated into an easy-to-use assay, we used three consensus DDR-related genes (*CXCL10*, *IDO1*, and *IFI44L*) from the 44- and 9-gene DDIR signatures to generate a novel DRIA signature. As predicted, the DRIA signature had a strong correlation with two original DDIR signatures and effectively classified patients into different subsets with distinct tumor immune microenvironment characteristics. Moreover, the level of *CXCL10*, *IDO1*, and *IFI44L* gene expression was shown to be increased in DDR-deficient tumors compared to DDR-proficient tumors, which indicated the need for three genes within the DDIR characteristics. In agreement with our results, Parkes et al.<sup>23</sup> concluded that DDR-deficient tumor cells express significantly higher *CXCL10* than DDR-proficient tumor cells. Vidotto et al.<sup>39</sup> reported that the increased expression of *IDO1* driven by DDR genetic deficiency modulates the response to immunotherapy. These data provided a theoretical basis for the DRIA signature to predict the response to ICI therapy.





Figure 6 Continued



**Figure 6** Validation of the prognostic value of DRIA in ICI-treated pan-cancer cohorts. (A-C) Kaplan–Meier plots of PFS segregated by the DRIA score in PUCH melanoma (A), Gide19 melanoma (B), and Pender21 pan-cancer cohorts (C). (D-F) Multivariate Cox regression analyses of DRIA and clinicopathologic factors associated with PFS in PUCH melanoma (D), Gide19 melanoma (E), and Pender21 pan-cancer cohorts (F). DRIA, DNA damage response-related immune activation; ICIs, immune checkpoint inhibitors; PFS, progression-free survival; HR, hazard ratio; CI, confidence interval.

The multi-cohort analysis and validation showed that the DRIA signature is predictive of clinical response and survival benefit following ICI therapy in patients with GI cancer and pan-cancer. The predictive accuracy of DRIA was superior to PD-L1, TMB, EBV, and MSI in patients with GI cancer. We also showed that the DRIA predictive performance was essentially equivalent to the two original DDIR signatures. In fact, these data fully demonstrated that the DRIA signature might serve

as a promising predictor for ICI therapy decisions. Further investigations are warranted to develop the detection system of three genes using a PCR array or liquid biopsy and conduct prospective evaluation of the DRIA signature. In addition, we showed that the pairwise combination of DRIA with PD-L1, TMB, MSI, and EBV had improved patient stratification for ICI therapy. These data suggested that the DRIA signature can also be used to complement these current biomarkers

to select potential responders for ICI therapy. Similarly, our study included one MSS GI cancer (Parikh22) cohort and successfully predicted three patients who responded to ICI-based combination therapy using the DRIA signature. Other studies have confirmed the presence of DDR pathway aberrations in patients with MSS colorectal cancer<sup>40,41</sup>. Further investigations are needed to extensively evaluate the predictive value of DRIA for ICI-based combination therapy in additional cohorts.

This retrospective study had several limitations, including unresolved concerns and potential perspectives. First, the DRIA signature was assessed based on endoscopic or needle biopsy specimens with limited tumor clonality. A high level of intra-tumoral heterogeneity is associated with the response to ICI therapy<sup>42,43</sup>, thereby indicating the limitation of a single biopsy in the development of a predictive biomarker. This limitation can be partially resolved by the pooling of biopsy fragments from multiple sites within the tumor. Second, several factors that may have inevitably produced bias, such as the sequencing platform used, line of treatment, ICI therapeutic regimen used, and the limited types of solid tumors. Moreover, although one real-world pan-cancer cohort was included in this study to validate the predictive value of DRIA, the small sample sizes weakened the accuracy of the results. Of note, however, results were validated in multiple cohorts spanning Asian, European, American, and Australian populations. Considering the limitations and concerns mentioned above, prospective multicenter clinical trials with a larger number of patients with multiple solid tumors are warranted to assess the predictive value of DRIA. Furthermore, comprehensive analyses of the relationship between DRIA and known clinical biomarkers based on a standardized testing system may provide additional information for guiding clinical application of ICI therapy.

## Conclusions

In summary, we constructed an individualized DDR-based signature (DRIA) that identified potential patients with improved response rates and survival outcomes in the context of ICI therapy. Our study also demonstrated that this signature has superior predictive power for the clinical benefit of ICI therapy compared to known clinical biomarkers, including PD-L1 TMB, MSI, and EBV. Therefore, DRIA may potentially act as a novel predictive marker for ICI therapy. More in depth corollary studies are warranted in prospective clinical trials.

## Grant support

This study was supported by the National Natural Science Foundation of China (Grant Nos. 81972761 and 82202837) and the National Key R&D Program of China (Grant Nos. 2016YFC1303200 and 2022YFC2505100).

## Conflict of interest statement

No potential conflicts of interest are disclosed.

## Author contributions

Conceived and designed the analysis: Yongzhan Nie, Yan Kong. Collected the data: Junya Yan, Shibo Wang, Jing Zhang, Qiangqiang Yuan, Xianchun Gao, Nannan Zhang, Haohao Zhang, Kun Liu, Jun Yu, Linbin Lu, Hui Liu, Xiaoliang Gao, Sheng Zhao, Wenyao Zhang, Abudurousuli Reyila, Yu Qi, Qiujin Zhang.

Provided the materials or patients: Yongzhan Nie, Yan Kong, Yan Pan, Shundong Cang, Yuanyuan Lu, Yanglin Pan.

Analyzed and interpreted the data: Yongzhan Nie, Yan Kong, Junya Yan, Shibo Wang, Jing Zhang.

Wrote the manuscript: Junya Yan.

## Data availability statement

The results shown here are in part based on data generated by The Cancer Genome Atlas project (TCGA, <http://cancergenome.nih.gov/>) and data available in published articles<sup>10,29-32</sup>. The raw sequence data from PUCH dataset reported in this paper have been deposited in the Genome Sequence Archive in National Genomics Data Center, Beijing Institute of Genomics (China National Center for Bioinformatics), Chinese Academy of Sciences, under accession number HRA000524 that are publicly accessible at <http://bigd.big.ac.cn/gsa-human>.

## References

1. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global Cancer Statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2021; 71: 209-49.
2. Javle M, Hsueh CT. Recent advances in gastrointestinal oncology – updates and insights from the 2009 annual meeting of the American Society of Clinical Oncology. *J Hematol Oncol.* 2010; 3: 11.

3. Wang Y, Wang M, Wu HX, Xu RH. Advancing to the era of cancer immunotherapy. *Cancer Commun (Lond)*. 2021; 41: 803-29.
4. Sun JM, Shen L, Shah MA, Enzinger P, Adenis A, Doi T, et al. Pembrolizumab plus chemotherapy versus chemotherapy alone for first-line treatment of advanced oesophageal cancer (KEYNOTE-590): a randomised, placebo-controlled, phase 3 study. *Lancet*. 2021; 398: 759-71.
5. Diaz LA Jr, Shiu KK, Kim TW, Jensen BV, Jensen LH, Punt C, et al. Pembrolizumab versus chemotherapy for microsatellite instability-high or mismatch repair-deficient metastatic colorectal cancer (KEYNOTE-177): final analysis of a randomised, open-label, phase 3 study. *Lancet Oncol*. 2022; 23: 659-70.
6. Wang ZX, Pan YQ, Li X, Tsubata T, Xu RH. Immunotherapy in gastrointestinal cancers: advances, challenges, and countermeasures. *Sci Bull (Beijing)*. 2023; 68: 763-6.
7. Shitara K, Özgüroğlu M, Bang YJ, Di Bartolomeo M, Mandalà M, Ryu MH, et al. Pembrolizumab versus paclitaxel for previously treated, advanced gastric or gastro-oesophageal junction cancer (KEYNOTE-061): a randomised, open-label, controlled, phase 3 trial. *Lancet*. 2018; 392: 123-33.
8. Wang F, Wei XL, Wang FH, Xu N, Shen L, Dai GH, et al. Safety, efficacy and tumor mutational burden as a biomarker of overall survival benefit in chemo-refractory gastric cancer treated with toripalimab, a PD-1 antibody in phase Ib/II clinical trial NCT02915432. *Ann Oncol*. 2019; 30: 1479-86.
9. Le DT, Uram JN, Wang H, Bartlett BR, Kemberling H, Eyring AD, et al. PD-1 blockade in tumors with mismatch-repair deficiency. *N Engl J Med*. 2015; 372: 2509-20.
10. Kim ST, Cristescu R, Bass AJ, Kim KM, Odegaard JI, Kim K, et al. Comprehensive molecular characterization of clinical responses to PD-1 inhibition in metastatic gastric cancer. *Nat Med*. 2018; 24: 1449-58.
11. Venderbosch S, Nagtegaal ID, Maughan TS, Smith CG, Cheadle JP, Fisher D, et al. Mismatch repair status and BRAF mutation status in metastatic colorectal cancer patients: a pooled analysis of the CAIRO, CAIRO2, COIN, and FOCUS studies. *Clin Cancer Res*. 2014; 20: 5322-30.
12. Kojima T, Shah MA, Muro K, Francois E, Adenis A, Hsu CH, et al. Randomized Phase III KEYNOTE-181 Study of pembrolizumab versus chemotherapy in advanced esophageal cancer. *J Clin Oncol*. 2020; 38: 4138-48.
13. Ott PA, Bang YJ, Piha-Paul SA, Razak A, Bannouna J, Soria JC, et al. T-cell-inflamed gene-expression profile, programmed death ligand 1 expression, and tumor mutational burden predict efficacy in patients treated with pembrolizumab across 20 cancers: KEYNOTE-028. *J Clin Oncol*. 2019; 37: 318-27.
14. Samstein RM, Lee CH, Shoushtari AN, Hellmann MD, Shen R, Janjigian YY, et al. Tumor mutational load predicts survival after immunotherapy across multiple cancer types. *Nat Genet*. 2019; 51: 202-6.
15. Lawrence MS, Stojanov P, Polak P, Kryukov GV, Cibulskis K, Sivachenko A, et al. Mutational heterogeneity in cancer and the search for new cancer-associated genes. *Nature*. 2013; 499: 214-8.
16. Chabanon RM, Rouanne M, Lord CJ, Soria JC, Pasero P, Postel-Vinay S. Targeting the DNA damage response in immuno-oncology: developments and opportunities. *Nat Rev Cancer*. 2021; 21: 701-17.
17. Pujade-Lauraine E, Ledermann JA, Selle F, GebSKI V, Penson RT, Oza AM, et al. Olaparib tablets as maintenance therapy in patients with platinum-sensitive, relapsed ovarian cancer and a BRCA1/2 mutation (SOLO2/ENGOT-Ov21): a double-blind, randomised, placebo-controlled, phase 3 trial. *Lancet Oncol*. 2017; 18: 1274-84.
18. Hugo W, Zaretsky JM, Sun L, Song C, Moreno BH, Hu-Lieskovan S, et al. Genomic and transcriptomic features of response to anti-PD-1 therapy in metastatic melanoma. *Cell*. 2016; 165: 35-44.
19. Teo MY, Seier K, Ostrovnaya I, Regazzi AM, Kania BE, Moran MM, et al. Alterations in DNA damage response and repair genes as potential marker of clinical benefit from PD-1/PD-L1 blockade in advanced urothelial cancers. *J Clin Oncol*. 2018; 36: 1685-94.
20. Wang Y, Jiao X, Li S, Chen H, Wei X, Liu C, et al. Alterations in DNA damage response and repair genes as potential biomarkers for immune checkpoint blockade in gastrointestinal cancer. *Cancer Biol Med*. 2021; 19: 1139-49.
21. Huang R, Zhou PK. DNA damage repair: historical perspectives, mechanistic pathways and clinical translation for targeted cancer therapy. *Signal Transduct Target Ther*. 2021; 6: 254.
22. Mulligan JM, Hill LA, Deharo S, Irwin G, Boyle D, Keating KE, et al. Identification and validation of an anthracycline/cyclophosphamide-based chemotherapy response assay in breast cancer. *J Natl Cancer Inst*. 2014; 106: djt335.
23. Parkes EE, Walker SM, Taggart LE, McCabe N, Knight LA, Wilkinson R, et al. Activation of STING-dependent innate immune signaling by S-phase-specific DNA damage in breast cancer. *J Natl Cancer Inst*. 2017; 109: djw199.
24. Sharma P, Barlow WE, Godwin AK, Parkes EE, Knight LA, Walker SM, et al. Validation of the DNA damage immune response signature in patients with triple-negative breast cancer from the SWOG 9313c Trial. *J Clin Oncol*. 2019; 37: 3484-92.
25. Parkes EE, Savage KI, Lioe T, Boyd C, Halliday S, Walker SM, et al. Activation of a cGAS-STING-mediated immune response predicts response to neoadjuvant chemotherapy in early breast cancer. *Br J Cancer*. 2022; 126: 247-58.
26. Turkington RC, Knight LA, Blayney JK, Secrier M, Douglas R, Parkes EE, et al. Immune activation by DNA damage predicts response to chemotherapy and survival in oesophageal adenocarcinoma. *Gut*. 2019; 68: 1918-27.
27. Malla SB, Fisher DJ, Domingo E, Blake A, Hassanieh S, Redmond KL, et al. In-depth clinical and biological exploration of DNA damage immune response as a biomarker for oxaliplatin use in colorectal cancer. *Clin Cancer Res*. 2021; 27: 288-300.
28. Cui C, Xu C, Yang W, Chi Z, Sheng X, Si L, et al. Ratio of the interferon- $\gamma$  signature to the immunosuppression signature predicts anti-PD-1 therapy response in melanoma. *NPJ Genom Med*. 2021; 6: 7.
29. Parikh AR, Szabolcs A, Allen JN, Clark JW, Wo JY, Raabe M, et al. Radiation therapy enhances immunotherapy response in microsatellite stable colorectal and pancreatic adenocarcinoma in a phase II trial. *Nat Cancer*. 2021; 2: 1124-35.

30. Gide TN, Quek C, Menzies AM, Tasker AT, Shang P, Holst J, et al. Distinct immune cell populations define response to anti-PD-1 monotherapy and anti-PD-1/anti-CTLA-4 combined therapy. *Cancer Cell*. 2019; 35: 238-55.e6.
31. Mariathasan S, Turley SJ, Nickles D, Castiglioni A, Yuen K, Wang Y, et al. TGF $\beta$  attenuates tumour response to PD-L1 blockade by contributing to exclusion of T cells. *Nature*. 2018; 554: 544-8.
32. Pender A, Titmuss E, Pleasance ED, Fan KY, Pearson H, Brown SD, et al. Genome and transcriptome biomarkers of response to immune checkpoint inhibitors in advanced solid tumors. *Clin Cancer Res*. 2021; 27: 202-12.
33. Teo MY, Bambury RM, Zabor EC, Jordan E, Al-Ahmadie H, Boyd ME, et al. DNA damage response and repair gene alterations are associated with improved survival in patients with platinum-treated advanced urothelial carcinoma. *Clin Cancer Res*. 2017; 23: 3610-8.
34. You Z, Lv M, He X, Pan Y, Ge J, Hu X, et al. Homologous recombination repair gene mutations as a predictive biomarker for immunotherapy in patients with advanced melanoma. *Front Immunol*. 2022; 13: 871756.
35. Wang Z, Zhao J, Wang G, Zhang F, Zhang Z, Zhang F, et al. Comutations in DNA damage response pathways serve as potential biomarkers for immune checkpoint blockade. *Cancer Res*. 2018; 78: 6486-96.
36. Linger RJ, Kruk PA. BRCA1 16 years later: risk-associated BRCA1 mutations and their functional implications. *FEBS J*. 2010; 277: 3086-96.
37. Esteller M, Silva JM, Dominguez G, Bonilla F, Matias-Guiu X, Lerma E, et al. Promoter hypermethylation and BRCA1 inactivation in sporadic breast and ovarian tumors. *J Natl Cancer Inst*. 2000; 92: 564-9.
38. Turner N, Tutt A, Ashworth A. Hallmarks of 'BRCAness' in sporadic cancers. *Nat Rev Cancer*. 2004; 4: 814-9.
39. Vidotto T, Nersesian S, Graham C, Siemens DR, Koti M. DNA damage repair gene mutations and their association with tumor immune regulatory gene expression in muscle invasive bladder cancer subtypes. *J Immunother Cancer*. 2019; 7: 148.
40. Song Y, Huang J, Liang D, Hu Y, Mao B, Li Q, et al. DNA damage repair gene mutations are indicative of a favorable prognosis in colorectal cancer treated with immune checkpoint inhibitors. *Front Oncol*. 2020; 10: 549777.
41. Keenan BP, VAN Loon K, Khilnani AD, Fidelman N, Behr SC, Atreya CE, et al. Molecular and radiological features of microsatellite stable colorectal cancer cases with dramatic responses to immunotherapy. *Anticancer Res*. 2021; 41: 2985-92.
42. Hu J, Chen Z, Bao L, Zhou L, Hou Y, Liu L, et al. Single-cell transcriptome analysis reveals intratumoral heterogeneity in cCRCC, which results in different clinical outcomes. *Mol Ther*. 2020; 28: 1658-72.
43. Nicos M, Krawczyk P, Crosetto N, Milanowski J. The role of intratumor heterogeneity in the response of metastatic non-small cell lung cancer to immune checkpoint inhibitors. *Front Oncol*. 2020; 10: 569202.

**Cite this article as:** Yan J, Wang S, Zhang J, Yuan Q, Gao X, Zhang N, et al. DNA damage response-related immune activation signature predicts the response to immune checkpoint inhibitors: from gastrointestinal cancer analysis to pan-cancer validation. *Cancer Biol Med*. 2024; 21: 252-266. doi: 10.20892/j.issn.2095-3941.2023.0303